250-477-0322

WWW.VICTORIAPESTCONTROL.COM

Quality Service Since 1986

FLEA TREATMENT NOTIFICATION

Siphonaptera

|--|



Dear Owner/Tenant: On inspection or identification we have confirmed Flea activity within your unit or home. In order for us to perform the best treatment we can, we ask you to please complete the following:

- * All pets must be on a Veterinarian administered flea program such as Advantage or Revolution.
- * Please vacuum daily before and after treatment. This is to stimulate fleas to hatch from their pupal cocoons and contact treatments. It may cause temporary increases in adult fleas.
- * Focus cleaning and vacuuming on areas where pets rest, this is often an area of flea development. Dispose of vacuum bags in sealed bags.
- * **Remove** all belongings from floor areas, including under beds and couches.
- * Remove or cover any exposed food or utensils.
- * Cover all fish tanks and disconnect air filter pumps.

Fleas may be seen after treatment due to the presence of different life stages not affected by pesticides. Control will be achieved, but new adult fleas hatching from pupal cocoons may be found after treatment. Most fleas cannot lay functional eggs without a non-human animal host present.

BEFORE OR AFTER TREATMENT, ALL BEDDING AND PILLOWS SHOULD BE HOT LAUNDERED, OR PUT THROUGH THE HOT CYCLE OF THE DRYER FOR 20-30 MIN. DO NOT OVER PACK DRYERS.

PESTICIDE USE NOTICE

Precor (Insect Growth Regulator): Methoprene 1.2% PCP# 15255

Dragnet (Adulticide): Permethrin 0.5% PCP# 24175

Seclira WSG Dinotefuran 0.1% PCP#34816

	AI:
Amount:	PCP#



BC Poison Control Information Centre 1-800-567-8911

For the above products Health Canada recommends that all pets and occupants vacate until treatments have dried. Please do not enter treated areas for 2-4 hours during and after treatment. Please ventilate your unit (open windows) upon re-entry.

